Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

CERAPP: Collaborative Estrogen Receptor Activity Prediction Project

Kamel Mansouri, Ahmed Abdelaziz, Aleksandra Rybacka, Alessandra Roncaglioni, Alexander Tropsha, Alexandre Varnek, Alexey Zakharov, Andrew Worth, Ann M. Richard, Christopher M. Grulke, Daniela Trisciuzzi, Denis Fourches, Dragos Horvath, Emilio Benfenati, Eugene Muratov, Eva Bay Wedebye, Francesca Grisoni, Giuseppe F. Mangiatordi, Giuseppina M. Incisivo, Huixiao Hong, Hui W. Ng, Igor V. Tetko, Ilya Balabin, Jayaram Kancherla, Jie Shen, Julien Burton, Marc Nicklaus, Matteo Cassotti, Nikolai G. Nikolov, Orazio Nicolotti, Patrik L. Andersson, Qingda Zang, Regina Politi, Richard D. Beger, Roberto Todeschini, Ruili Huang, Sherif Farag, Sine A. Rosenberg, Svetoslav Slavov, Xin Hu, and Richard S. Judson

Table of Contents

Table S1: Information on participating groups

Table S2: Project plan and tasks accomplished in each step

Table S3: Statistics and scores of the categorical models of the different research groups in alphabetic order.

Table S4: Sensitivity and specificity of the different evaluation steps for categorical models

Table S5: statistics and scores of agonist, antagonist and binding continuous models based on the five potency classes defined using the reference chemicals.

Table S6: Participant single categorical models compared to agonist antagonist and binding categorical *consensus* predictions (as the "observed response") after the 4 correction rules were applied. It shows the number of predicted chemicals (out of the total 32k prediction set), the number of actives (out of the 4001 actives predicted by the consensus categorical model) and summary statistics based on the consensus categorical predictions.

Figure S1: Number of chemicals by positive concordance (agreement on actives between the included models) of the categorical binding models on the active compounds of the prediction set (32k). The positive concordance is the fraction of models predicting a certain chemical as active by the total number of models providing a prediction for that chemical.

Figure S2: Bar plot of the corrected categorical *consensus* for binding accuracy with variable number of literature sources from the evaluation set. The y –axis, depending on the color of the bar from the legend, represents the balanced accuracy, sensitivity or specificity.